Trockenfutter

Pellets

Strukturfutter

Heu

Buntfutter

Vergleiche

(siehe auch Nahrung, Verhalten, Futtermittelanalyse)

Rothacher *et al*., 2023[1)] verglichen handelsübliche Futtermittel mit Kenntnissen zur Ernährung und zum Fressverhalten von Kaninchen der Art *O. cuniculus* in natürlichen Lebensräumen.
Die Suche nach Primärliteratur über die quantitative Zusammensetzung natürlicher Nahrung und das Aktivitätsverhalten freilebender Kaninchen ergab folgende Ergebnisse (Tabelle 1–3).

Tabelle 1: Pflanzliche Zusammensetzung der Nahrung (nach [2)])

Einkeimblättrige (Gräser)	Zweikeimblättrige (Kräuter, krautige Pflanzen, Weidepflanzen)	Ein- und Zweikeimblättrige (inkl. nicht identifizierte Pflanzen)	Früchte	Saaten	Referenzen
16–86 %	10–83 %	89–100 %	–	0–11 %	[3)4)5)][6)][7)][8)][9)][10)]

Tabelle 2: Nährstoffzusammensetzung der Nahrung in g/kg Trockensubstanz (nach [11)])

Bedingung	Protein	NDF	Rohfaser	Referenzen
Spanien, ganzjährig	50–95	–	–	[12)]
Niederlande, ganzjährig	198	570	275	[13)]
Frankreich, Frühling/Sommer	120–200	531–610	250–300	[14)]

Tabelle 3: Dauer des Nahrungsaufnahmeverhaltens in Stunden (nach [15)])

Bedingung	Nahrungsaufnahme/-suche	Referenzen
Frei lebend	3,8–7,0	[16)]
Heimtier, nur Heu	10,2	[17)]
Heimtier, nur Müsli-Mischung	2,5	[18)]
Versuchstier, Alleinfutter	2,2	[19)]

Anschließend wurden 134 Mischfuttermittel identifiziert, die im Zeitraum 1987–2020 für Heim- oder Nutzkaninchen kommerziell erhältlich waren (Allein- oder Ergänzungsfutter; pelletiert, extrudiert oder „Müsli“), und deren Zusammensetzung sowie die Fütterungsempfehlungen der Hersteller kritisch beleuchtet.

---

2 1 203

[1)](#), [2)](#), [11)](#), [15)](#)
Rothacher, M., Hatt, J. M., & Clauss, M. (2023). A comparison of commercially available feeds for rabbits, guinea pigs, chinchillas and degus with evidence of their diet and feeding behaviour in natural habitats. Schweizer Archiv fur Tierheilkunde, 165(11), 726-736.

[3)](#)
Alves, J. A. B., Vingada, J., & Rodrigues, P. (2006). The wild rabbit (Oryctolagus cuniculus) diet on a sand dune area in central Portugal: a contribution towards management. Wildlife Biology in Practice, 2(2), 63-71.

[4)](#)
Bonino, N., & Borrelli, L. (2006). Variación estacional en la dieta del conejo silvestre europeo (Oryctolagus cuniculus) en la región andina de Neuquén, Argentina. Ecología austral, 16(1), 7-13.

[5)](#)
Katona, K., Bíró, Z., Hahn, I., Kertész, M., & Altbacker, V. (2004). Competition between European hare and European rabbit in a lowland area, Hungary: a long-term ecological study in the period of rabbit extinction. Folia Zoologica, 53(3), 255-268.

[6)](#)
Marques, C., & Mathias, M. L. (2001). The diet of the European wild rabbit, Oryctolagus cuniculus (L.), on different coastal habitats of Central Portugal. Mammalia; 65: 437–449.

[7)](#)
Martin, G. R., Twigg, L. E., & Zampichelli, L. (2007). Seasonal changes in the diet of the European rabbit (Oryctolagus cuniculus) from three different Mediterranean habitats in south-western Australia. Wildlife Research, 34(1), 25-42.

[8)](#)
Robley, A. J., Short, J., & Bradley, S. (2001). Dietary overlap between the burrowing bettong (Bettongia lesueur) and the European rabbit (Oryctolagus cuniculus) in semi-arid coastal Western Australia. Wildlife Research, 28(4), 341-349.

[9)](#)
Rogers, P. M., Soriguer, R. C., & Arthur, C. P. (1994). The rabbit in continental Europe. In: The European rabbit (eds Thompson, H. V. & King, C. M.). Oxford University Press. 22-63.

[10)](#)
Wolfe, A., Whelan, J., & Hayden, T. J. (1996). Dietary overlap between the Irish mountain hare Lepus timidus hibernicus and the rabbit Oryctolagus cuniculus on coastal grassland. In: Biology and Environment. Proceedings of the Royal Irish Academy. 89-95.

[12)](#)
Lombardi, L., Fernández, N., & Moreno, S. (2007). Habitat use and spatial behaviour in the European rabbit in three Mediterranean environments. Basic and Applied Ecology, 8(5), 453-463.

[13)](#)
Deinum, B., & Wallage-Drees, J. M. (1985). Quality of the diet selected by wild rabbits (Oryctolagus cuniculus) in autumn and winter. Netherlands Journal of Zoology, 36(4), 438-448.

[14)](#)
Thompson, H. V., & King, C. M. (1994). The European rabbit, the history and biology of a successful colonizer. Oxford Science Publications, 1994.

[16)](#)
Mykytowycz, R., & Rowley, I. (1958). Continuous observations of the activity of the wild rabbit, Oryctolagus cuniculus, during 24 hour periods. CSIRO Wildlife Research, 3(1), 26-31.

[17] [18], Prebble, J. L., Langford, F. M., Shaw, D. J., & Meredith, A. L. (2015). The effect of four different feeding regimes on rabbit behaviour. Applied Animal Behaviour Science, 169, 86-92.
[19] Olivas, I., Rodriguez-Latorre, A., Estellés, F., Calvet, S., & Villagrá, A. (2013). Assessment of activity patterns of growing rabbits in a flux-controlled chamber. World Rabbit Science, 21(2), 107-110.

From:
https://wikikanin.de/ - **Wikikanin**

Permanent link:
https://wikikanin.de/doku.php?id=futtermittel:trockenfutter&rev=1779821621

Last update: **2026/05/26 20:53**